

# No Lineal 2018

International Conference on Nonlinear Mathematics  
and Physics



Málaga, June 12th-15th, 2018  
<http://nonlinear18.uma.es>



Book of Abstracts

Book of Abstracts  
**International Conference on Nonlinear  
Mathematics and Physics Malaga 2018**

Editors

**Francisco R. Villatoro, Francisco Rus, Julio Garralon  
and Francisca Martin-Vergara**

**ISBN: 978-84-09-03289-1 (No comercial)**

This book is available from

**<http://nonlinear18.uma.es/book-of-abstracts/>**


Supported by



EphemeCH research group  
TIN2014-56494-C4-1..4-P



Academia Malagueña de Ciencias



Programa Doctorado UMA de Ingeniería Mecánica y Eficiencia Energética



# Table of Contents

<b>EDITORIAL COMMENTS</b>	<b>I</b>
<b>Committees</b>	<b>II</b>
<b>Program</b>	<b>III</b>
<b>Plenary Talks</b>	<b>IV</b>
<b>Divulgative Talk</b>	<b>V</b>
<b>Oral communications</b>	<b>VI</b>
<b>On a model for internal waves in a rotating fluid</b> Duran, Angel	<b>1</b>
<b>Center cyclicity on center manifolds of polynomial families of vector fields</b> Maza, Susanna	<b>2</b>
<b>Leaky surface wave propagation for mass loading biosensors: A numerical study</b> Craig, Nathan and Grigg, Harriet	<b>3</b>
<b>Bautin-type analysis in 1-dimensional averaged theory for periodic differential equations</b> Garcia, Isaac A.	<b>5</b>
<b>Periodic solutions for an scalar differential equation modelling optical conveyor belts</b> Valero, Jose and Carretero, Luis	<b>6</b>
<b>Information flow in a two-scale stochastic dynamical system using ordinal symbolic analysis</b> Rosa, Santiago, Wiedmad Santiago, Jan Van Leeuwen, Peter, Lucivi, Magdalena and Pulido, Manuel	<b>7</b>
<b>Multistability in the dynamics of bubble contrast agents</b> Garashchuk, Ivan R., Sinelshchikov, Cimitry I and Kudryashov, Nikolay A.	<b>8</b>
<b>Robustness of dynamically gradient multivalued dynamical systems</b> Caballero, Ruben, Carvalho, Alexandre N., Marin-Rubio, Pedro and Valero, Jose	<b>10</b>
<b>Global connections near a triple-zero singularity</b> Algaba, A., Dominguez-Moreno, M.C. and Merino, M.	<b>11</b>
<b>Computational methods for compactons in the Rosenau-Hyman equation</b> Rus, Francisco and Villatoro, Francisco	<b>12</b>
<b>On the stability of positive periodic solutions in MEMs via lower and upper solutions</b> Nuñez, Daniel	<b>13</b>
<b>Finite difference methods based on time-linearization for the GRLW equation</b> Garcia-Lopez, C.M. and Ramos, J.I.	<b>14</b>
<b>Time-diffracting beams in self-focusing Kerr media: self trapped propagation below the critical power</b> Porras, Miguel A.	<b>16</b>

<b>A necessary condition to have Weierstrass integrability</b> Gine, Jaume and Llibre Jaume	17
<b>Analytic integrability of planar systems with non null linear part</b> Algaba, Antonio, Diaz, Maria, Garcia, Cristobal and Ginev, Jaume	18
<b>Supressing chaos in damped driven systems by incommensurate excitations</b> Chacon, R., Miralles, J. J., Martinez, J. A. and Balibrea, F.	19
<b>The role of discrete breathers in establishing the Fourier heat transport in <math>\phi^4</math> chain</b> Dmitriev, Sergey V. and Xiong, Daxing	21
<b>Orbital-Reversibility for vector fields. Application to center problem</b> Algaba, A., Checa, I. and Garcia, C.	22
<b>Generalising the BDS test to check i.i.d. in high-frecuency finantial time series</b> Sandubete, Julio E. and Escot, Lorenzo	24
<b>Punchets: nonlinear transport in Hamiltonian pump-ratchet hybrids</b> Dittrich, Thomas and Medina Sanchez, Nicolas	25
<b>Padé numerical schemes for the sine-Gordon equation</b> Martin-Vergara, F. and Villatoro, F. R.	26
<b>Localized vibrational modes in metals and alloys</b> Korznikova, Elena A.	27
POSTER	
<b>Long-lived vortex soliton arrays nested in flat-top nonlinear Bessel beams</b> Porras, Miguel A., Ramos, Francisco and Garcia-Riquelme, Jose L.	28
<b>Participants</b>	VII

The 11th International Conference on Nonlinear Mathematics and Physics (NoLineal 2018) will be held in Málaga (Spain), 12-15th June, 2018. This series of conferences were previously held in Ávila (1997), Almagro (2000), Cuenca (2002), Toledo (2004), Ciudad Real (2007), Barcelona (2008), Cartagena (2010), Zaragoza (2012), Badajoz (2014) and Sevilla (2016).

The aim of this conference is to offer senior and young researchers of different areas related to nonlinear sciences, such as Physics, Mathematics, Biology, Economics, Social Sciences, etc, the possibility to share their latest results in this interdisciplinary meeting.

## Scientific Committee

Francisco Balibrea Gallego (Departamento de Matemáticas, Universidad de Murcia, Spain)  
Ricardo Chacón (Universidad de Extremadura, Spain)  
Leonor Cruzeiro (Universidade do Algarve, Portugal)  
Faustino Palmero Acebedo (Universidad de Sevilla, Spain)  
Avadh B. Saxena (Center for Nonlinear Studies, Los Alamos National Laboratory, USA)  
Miguel Ángel Rodríguez (Instituto de Física de Cantabria, Universidad de Cantabria, Spain)  
Josep J. Masdemont (Universitat Politècnica de Catalunya, Spain)  
Gustavo Deco (Universitat Pompeu Fabra, Spain)  
Jordi Caballe (Univesitat Autònoma de Barcelona, Spain)  
Luis Martínez Salamero (Universitat Rovira i Virgili, Spain)  
Ángel Jorba (Universitat de Barcelona, Spain)  
Ignacio J. Turias (Universidad de Cádiz, Spain)  
Juan Pablo Neirotti (University of Aston, United Kingdom)  
Laura Rebollo-Neira (University of Aston, United Kingdom)  
Juan Perotti (Universidad de Córdoba, Argentina)  
Manuel Pulido (Universidad del Litoral, Argentina)  
Ricardo Carretero (San Diego State University, USA)  
Humberto Michinel (Universidad de Vigo, Spain)

## Local Scientific Committee

Carlos Cotta Porras (Universidad de Málaga, Spain)  
Antonio José Fernández Leiva (Universidad de Málaga, Spain)  
Leonardo Franco (Universidad de Málaga, Spain)  
Carmen María García López (Universidad de Málaga, Spain)  
Julio Garralón Ruiz (Universidad de Málaga, Spain)  
José Manuel Jerez Aragonés (Universidad de Málaga, Spain)  
Francisco Rus Mansilla (Universidad de Málaga, Spain)  
Miguel Atencia (Universidad de Málaga, Spain)

## Local Organizing Committee

Julio Garralón Ruiz (Universidad de Málaga, Spain)  
Francisca Martin-Vergara (Universidad de Málaga, Spain) Francisco Rus  
Mansilla (Universidad de Málaga, Spain) Francisco R. Villatoro  
Machuca (Universidad de Málaga, Spain)



# XI International Conference on Nonlinear Mathematics and Physics

Malaga, June 12th – 15th, 2018

Escuela de Ingenierías Industriales, Sala de Grados A  
(Ampliación del Campus de Teatinos)

## Program

Tuesday, 12th June

---

- 09:00 - 10:15 Registration  
10:15 - 10:30 Presentation  
10:30 - 11:30 Plenary session: B. Malomed  
Multidimensional solitons  
11:30 - 12:00 Coffee break  
12:00 - 13:00 Plenary session: P. Rosenau  
Compactons in Action: When, Where and How  
13:30 - 15:30 Lunch  
15:30 - 17:10 Afternoon scientific sessions: lecture room  
15:30 A. Durán, 15:55 S. Maza, 16:20 N. Craig, 16:45 I.A. García

Wednesday, 13th June

---

- 09:00 - 10:00 Plenary session: R. Carretero  
Vortices and vortex rings in quantum superfluids: a quasi-particle approach  
10:00 - 11:00 Plenary session: J.F.R. Archilla  
Nonlinear excitations in solids, observation and consequences  
11:00 - 11:30 Coffee break  
11:30 - 13:35 Morning scientific sessions: lecture room  
11:30 J. Valero, 11:55 M. Pulido, 12:20 D. Sinelshchikov, 12:45 R. Caballero, 13:10 M.C. Domínguez  
13:40 - 15:30 Lunch  
15:30 - 17:10 Afternoon scientific sessions: lecture room  
15:30 F. R. Villatoro, 15:55 D. Núñez, 16:20 C.M. García-López, 16:45 M.A. Porras  
18:30 Panoramic tour (downtown)

Thursday, 14th June

---

- 09:00 - 10:00 Plenary session: Y. Kosevich  
Phase dynamics of nonlinear localized excitations in two weakly coupled anharmonic chains and analogy with coherent tunneling of quantum objects  
10:00 - 11:00 Plenary session: Y. Kartashov  
Nonlinear phenomena in polariton topological insulators  
11:00 - 11:30 Coffee break  
11:30 - 13:35 Morning scientific sessions: lecture room  
11:30 J. Giné, 11:55 M. Díaz García, 12:20 J.J. Miralles, 12:45 S. V. Dmitriev, 13:10 I. Checa  
13:40 - 15:30 Lunch  
15:30 - 17:10 Afternoon scientific sessions: lecture room  
15:30 J.E. Sandubete, 15:55 T. Dittrich, 16:20 F. Martín-Vergara, 16:45 E. Korznikova  
19:00 – 20:00 Popular talk: Manuel Pulido (Rectorate building, downtown)  
¿Es reversible el cambio climático antropogénico? Impacto del “agujero de ozono” en el clima del hemisferio sur  
20:30 Conference dinner (downtown)

Friday, 15th June

---

- 09:00 - 10:00 Plenary session: J. I. Ramos  
Superdiffusive Burgers-Taylor's kinks and nonlinear dispersive shock waves  
10:00 - 11:00 Plenary session: J. C. Losada  
From 2D to 3D in LiCN: A brief history of Nonlinear Dynamics of a Hamiltonian Molecular System  
11:00 Closing coffee



## Plenary Talks

## **Divulgative Talk**

## Oral communications

# On a model for internal waves in a rotating fluid

ANGEL DURAN

*Department of Applied Mathematics, University of Valladolid*  
angel@mac.uva.es

## ABSTRACT

The purpose of this talk is to bring up some aspects on a model for the propagation of internal waves in a rotating fluid. This is based on incorporating rotational effects in the problem for internal waves described by the Benjamin equation, [2, 3, 1], or, alternatively, on including capillary forces in the Ostrovsky equation, [4, 5], for wave dynamics in a rotating fluid. Preliminary results concerning the regime of validity, the well-posedness of the initial-value problem and the existence of solitary-wave solutions will be presented.

## References

- [1] J.P. Albert, J.L. Bona, J.M. Restrepo, Solitary-wave solutions of the Benjamin equation, *SIAM J. Appl. Math.* 59 (1999) 2139-2161.
- [2] T.B. Benjamin, A new kind of solitary wave, *J. Fluid Mech.* 245 (1992) 401-411.
- [3] T.B. Benjamin, Solitary and periodic waves of a new kind, *Phil. Trans. R. Soc. A* 354 (1996) 1775-1806.
- [4] R. H. Grimshaw, Evolution equations for weakly nonlinear, long internal waves in a rotating fluid, *Stud. Appl. Math.*, 73 (1985) 1-33.
- [5] L. A. Ostrovsky, Nonlinear internal waves in a rotating ocean, *Okeanologia*, 18 (1978) 181-191.

# Center cyclicity on center manifolds of polynomial families of vector fields

SUSANNA MAZA

*Departament de Matemàtica, Universitat de Lleida,  
Avda. Jaume II, 69, 25001 Lleida, Spain  
smaza@matematica.udl.cat*

## ABSTRACT

*We consider polynomial families of ordinary differential equations on  $\mathbb{R}^3$ , parametrized by the admissible coefficients, for which there is an isolated singularity at which the linear part of the system has one non-zero real and two purely imaginary eigenvalues. We derive theorems that bound the maximum number of limit cycles within the center manifold that can bifurcate, under arbitrarily small perturbation of the coefficients, from any center at the singularity. The bounds are global in that they apply to the system corresponding to any point on any irreducible component of the center variety in the space of parameters. We also derive theorems for such bounds when attention is confined to a single irreducible component of the center variety. Finally, we apply the results obtained to get the cyclicity of the center located at the origin of the well-known Lorenz system. Moreover, we extend the results obtained in [2] about the center cyclicity in the Moon-Rand family. The talk is based on the work [1].*

## References

- [1] I.A. GARCÍA, & S. MAZA, D. SHAFER, *Cyclicity of polynomial nondegenerate centers on center manifolds*, Preprint.
- [2] A. MAHDI, V. G. ROMANOVSKI, AND D. S. SHAFER, *Stability and periodic oscillations in the Moon-Rand systems*, *Nonlinear Anal. Real World Appl.* **14** (2013), 294—313.

# Leaky Surface Wave Propagation for Mass Loading Biosensors: A Numerical Study

NATHAN CRAIG<sup>1</sup> AND HARRIET GRIGG<sup>2</sup>

<sup>1</sup>*School of Engineering, Newcastle University, Newcastle Upon Tyne, UK.  
n.p.craig@newcastle.ac.uk*

<sup>2</sup>*School of Engineering, Newcastle University, Newcastle Upon Tyne, UK.  
harriet.grigg@newcastle.ac.uk*

## ABSTRACT

*Surface acoustic wave (SAW) devices are a promising candidate for Point-of-Care (PoC) testing [1]. These devices relate the mass loading of an analyte to a measurable shift in elastic wave properties. The properties are poorly understood outside of standard configurations such as ST-cut Quartz. Presented is the numerical study into the existence of plane waves in multilayered anisotropic structures, developed based on the Stroh formulation [2]. The effect of dissimilar anisotropies on the existence of leaky and non-leaky SAW modes is explored in a layered half-space model. A second model with an additional layer of viscoelastic material is used to study these waves with change in concentration of analyte. The slowness surfaces are presented for both cases; these are required for designing non-circular annular interdigital transducers [3] for novel circular resonators [4]. The shift in frequency due to changes in analyte concentration is investigated for particular propagation angles and velocity ranges. These cases are at transition states in which the propagating bulk waves allow leaky modes to exist. The differing symmetries of the anisotropic layers allow these leaky modes to come into existence at particular directions governed by the combination of bulk wave solutions between each layer. The structure of these waves suggest a mode of operation based on the behaviour at transition with the possible creation of discontinuities.*

## References

- [1] S. K. Vashist, P. B. Luppa, L. Y. Yeo, A. Ozcan, and J. H. T. Luong, “Emerging Technologies for Next Generation Point-of-Care Testing, Trends in Biotechnology,” Vol. 33, No. 11, pp. 692-705, 2015.
- [2] K. Tanuma, “Stroh Formulation and Rayleigh Waves,” Journal of Elasticity Vol. 89, No. 1-3, pp. 5-154, 2007.
- [3] V. Laude, K. Kokkonen, and S. Benchabane, “Characterization of surface acoustic wave focusing by an annular interdigital transducer,” in Ultrasonics Symposium IEEE, Rome, Italy, 2009.

- [4] J. S. Burdess, et al., “The theory of a trapped degenerate mode resonator,” The journal of the Acoustical Society of America, Vol. 141, No. 6, pp. 4622 - 4632, 2017.

# Bautin-type analysis in 1-dimensional averaging theory for periodic differential equations

ISAAC A. GARCÍA

*Departament de Matemàtica, Universitat de Lleida,  
Avda. Jaume II, 69, 25001 Lleida, Spain  
garcia@matematica.udl.cat*

## ABSTRACT

*The classical averaging theory is a tool to study the dynamics of the periodic nonlinear differential systems, see [2]. We use averaging theory for computing  $T$ -periodic solutions of  $\lambda$ -families of analytic  $T$ -periodic ordinary differential equations in standard form defined on  $\mathbb{R}$  analyzing the displacement functions  $d(z, \lambda, \varepsilon)$  defined by these equations. Recall that  $d : \Omega \times \mathbb{R}^p \times I \rightarrow \Omega$  is a real analytic functions,  $\Omega$  is a bounded open subset of  $\mathbb{R}$ ,  $I \subset \mathbb{R}$  is an interval containing the origin,  $\lambda \in \mathbb{R}^p$  are parameters, and  $\varepsilon$  is a small parameter. We study the branching of the zero-set of  $d$  when the parameter  $\varepsilon$  varies following a Bautin-type analysis [1]. We call the coefficients in the Taylor expansion of  $d(z, \lambda, \varepsilon)$  in powers of  $\varepsilon$  the averaged functions. The main contribution consists in analyzing the role that have the multiple zeros  $z_0 \in \Omega$  of the first non-zero averaged function. The outcome is that we bound the maximum number of branches of isolated zeros that can bifurcate from each multiple zero  $z_0$  in two different ways depending on whether  $(z_0, \lambda)$  belongs or not to the analytic set defined by the real variety associated to the ideal generated by the averaged functions in the Noetheriang ring of all the real analytic functions at  $(z_0, \lambda)$ . Several examples illustrate our results and they are compared with the classical theory, branching theory and also under the light of singularity theory of smooth maps. The examples range from polynomial vector fields to Abel differential equations and perturbed linear centers. The talk is based on the work [3].*

## References

- [1] N.N. BAUTIN, *On the number of limit cycles which appear with the variations of the coefficients from an equilibrium point of focus or center type*, AMS Translations-Series 1, 5, 1962, 396–413 [Russian original: Math. Sbornik, 30, 1952, 181–1961].
- [2] J.A. SANDERS, F. VERHULST AND J. MURDOCK, *Averaging Methods in Nonlinear Dynamical Systems*, Second edition, Applied Mathematical Sciences 59, Springer, New York, 2007.
- [3] I.A. GARCÍA, J. LLIBRE & S. MAZA, *On the multiple zeros of a real analytic function with applications to the averaging theory of differential equations*. To appear in Nonlinearity.



# Periodic solutions for an scalar differential equation modelling optical conveyor belts

JOSÉ VALERO

*Centro de Investigación Operativa  
Universidad Miguel Hernández de Elche, 03202, Elche, Spain  
jvalero@umh.es*

LUIS CARRETERO

*Departamento de Ciencia de Materiales, Óptica y Tecnología Electrónica  
Universidad Miguel Hernández de Elche, 03202, Elche, Spain  
luis@dite.umh.es*

## ABSTRACT

*In this work, we study the existence of periodic solutions for a differential equation modelling an optical conveyor belt, where the gradient forces that act on the  $z$ -axially confined particles are obtained by means of the superposition of two temporally dephased counter propagating complex electromagnetic fields. The dynamic of particles in an axial optical conveyor belt can be modeled by the differential equation:*

$$z' = F_z(t, z), \quad (1)$$

where

$$F_z(t, z) = \frac{\partial V(t, z)}{\partial z},$$
$$V(t, z) = F_0 f(z) \cos\left(kz - \frac{bt}{2}\right)^2,$$

with  $F_0, k, b > 0$ .

*Using a general theorem from [1] on existence of solutions for boundary-value problems we obtain the existence of periodic solutions for the nonautonomous equation (1) in the following particular cases:*

- *Lorentzian axial region strength:  $f(z) = \frac{1}{1+(z/z_0)^2}$ ,  $z_0 > 0$ .*
- *Gaussian axial region strength  $f(z) = \exp(-2\frac{z^2}{z_0^2})$ ,  $z_0 > 0$ .*

## References

- [1] I. Kiguradze, B. Puza, On boundary value problems for functional differential equations, Mem. Differential Equations Math. Phys., **12** (1997), 106-113.

# Information flow in a two-scale stochastic dynamical system using ordinal symbolic analysis

SANTIAGO ROSA (1), SANTIAGO WIEDMANN (2), PETER JAN VAN LEEUWEN (3), MAGDALENA LUCINI (3,1), AND MANUEL PULIDO (3,1)

(1) *FaMAF, Universidad Nacional de Cordoba (Argentina)*, (2) *FaCENA, Universidad Nacional del Nordeste (Argentina)*, (3) *Department of Meteorology, University of Reading (UK)*  
m.a.pulido@reading.ac.uk

## ABSTRACT

Although the resolved dynamical equations of atmospheric or oceanic global numerical models are well established, the development and evaluation of parameterizations that represent subgrid-scale effects pose a big challenge [1]. Small-scale waves in the geophysical systems generate in a region, then propagate conservatively long distances to finally break and deposit their momentum far away from their generation location. Therefore, the system is characterized by strong nonlinear and remote interactions between the different scales. To mimic these interactions between subgrid-scale and large-scale dynamics, we use a proof-of-concept two-scale stochastic dynamical system based on Lorenz 1996 with long-range interactions between the two-scales. In this work, time series obtained from these systems are evaluated with information flow measures, i.e. mutual information and transfer entropy[2]. The measures are obtained via ordinal symbolic analysis using the Bandt-Pompe[3] symbolic data reduction in the signals. By comparing different experiments, we show that even when the dynamics of the large-scale variables of the systems has a short-range correlations, small-scale variable may give place to long-range information flow. We evaluate the potencial of information measures as a tool for establishing the structure and functional dependencies to be represented by the subgrid-scale parameterizations. .

## References

- [1] Pulido M. and O. Rosso, 2017. Model selection: Using information measures from ordinal symbolic analysis to select model sub-grid scale parameterizations. *J. Atmos. Sci.*, **74**, 3253–3269 doi: 10.1175/JAS-D-16-0340.1
- [2] Staniek, M. and Lehnertz, K., 2008. Symbolic transfer entropy. *Phys. rev. lett.*, **100**, p.158101.
- [3] Bandt, C. and Pompe, B., 2002. Permutation entropy: a natural complexity measure for time series. *Phys. rev. lett.*, **88**, p.174102.

# Multistability in the dynamics of bubble contrast agents

IVAN R. GARASHCHUK, DMITRY I. SINELSHCHIKOV, NIKOLAY A. KUDRYASHOV

*Department of Applied Mathematics, National Research Nuclear University  
MEPhI, 31 Kashirskoe Shosse, 115409 Moscow, Russian Federation  
disine@gmail.com*

## ABSTRACT

In this talk we consider dynamical systems for the description of the oscillations of spherical gas bubbles in a liquid. Such systems can be used for modelling of ultrasound contrast agents, which are used for enhancing ultrasound visualization [1]. While there are a lot of different models of microbubble contrast agents and it is known that complex dynamics may occur in such models (see, e.g. [2, 3, 4]), only few of them were thoroughly studied from a dynamical systems point of view. In addition, the knowledge of the location in the parameter and initial conditions space of regions of chaotic and regular dynamics is important for applications since both types of the dynamics may be either beneficial or undesirable depending on a particular usage of microbubbles. Therefore, understanding of the nonlinear dynamics of microbubbles is an important problem.

Here we investigate two models of encapsulated gas bubbles dynamics. The first one takes into account the liquid viscosity and compressibility and bubble's shell. The shell term is chosen according to the de-Jong model [5]. In the second one we in addition take into consideration impact of an elastic wall near which a bubble oscillates. We show that in both of these models complex dynamics can occur. For example, there is a possibility of coexistence of two periodic attractors or periodic and chaotic ones. We study the dependence of dynamics type on the two control parameters that are magnitude and frequency of the external pressure field. We find regions of the parameters which correspond to regular, chaotic and multistable dynamics. To this aim we use several approaches, including the numerical continuation and the perpetual points methods (see, e.g. [6, 7]).

This work is supported by Russian Science Foundation, grant number 17-71-10241.

## References

- [1] L. Hoff, Acoustic characterization of contrast agents for medical ultrasound imaging, Springer, Berlin, 2001.
- [2] U. Parlitz, V. Englisch, C. Scheffczyk, W. Lauterborn, Bifurcation structure of bubble oscillators, J. Acoust. Soc. Am. 88 (1990) 1061–1077.

- [3] J.M. Carroll, M.L. Calvisi, L.K. Lauderbaugh, Dynamical analysis of the nonlinear response of ultrasound contrast agent microbubbles, *J. Acoust. Soc. Am.* 133 (2013) 2641–2649.
- [4] I. Garashchuk, D. Sinelshchikov, N. Kudryashov, Hidden Attractors in a Model of a Bubble Contrast Agent Oscillating Near an Elastic Wall. *EPJ Web of Conferences* 173 (2018) 06006.
- [5] A.A. Doinikov, A. Bouakaz, Review of shell models for contrast agent microbubbles, *IEEE Trans. Ultrason. Ferroelectr. Freq. Control.* 58 (2011) 981–993.
- [6] A. V. Borisov, A.O. Kazakov, I.R. Sataev, Spiral chaos in the nonholonomic model of a Chaplygin top, *Regul. Chaotic Dyn.* 21 (2016) 939–954.
- [7] D. Dudkowski, A. Prasad, S. Jafari, T. Kapitaniak, N.V. Kuznetsov, G.A. Leonov, G A., Hidden attractors in dynamical systems, *Physics Reports*, 637 (2016) 1–50.

# Robustness of dynamically gradient multivalued dynamical systems

RUBÉN CABALLERO<sup>1</sup>, ALEXANDRE N. CARVALHO<sup>2</sup>, PEDRO MARÍN-RUBIO<sup>3</sup>,  
AND JOSÉ VALERO<sup>1</sup>

<sup>1</sup>*Centro de Investigación Operativa, Universidad Miguel Hernández de Elche, Spain*

ruben.caballero@umh.es, jvalero@umh.es

<sup>2</sup>*Instituto de Ciências Matemáticas e de Computação, Universidade de São Paulo, Brazil*

andcarva@icmc.usp.br

<sup>3</sup>*Departamento de Ecuaciones Diferenciales y Análisis Numérico, Universidad de Sevilla, Spain*

pmr@us.es

## ABSTRACT

*In this work we show under suitable assumptions that a dynamically gradient multivalued semiflow is stable under perturbations. These results extend previous ones in the single-valued framework to the case where uniqueness of solution does not hold.*

*Afterwards, we consider a Chafee-Infante problem, where the equivalence of weak and strong solutions and the set of fixed points is analyzed. Also, the weak solutions of a particular family of Chafee-Infante problems approximating a differential inclusion studied in [1] generate a dynamically gradient multivalued semiflow with respect to suitable Morse sets. Moreover, we check that this family of Chafee-Infante equations verifies the hypotheses of the robustness theorem. In a more general context, describing the structure of global attractors is an open question on which focus efforts.*

## Acknowledgements

*This research was supported by the following projects: R. Caballero is a fellow of Programa de FPU del MECD, reference FPU15/03080; P. Marín-Rubio was supported by projects PHB2010-0002-PC (MECD-DGPU), MTM2015-63723-P (MINECO/FEDER, EU) and P12-FQM-1492 (Junta de Andalucía); and J. Valero by projects MTM2015-63723-P, MTM2016-74921-P (MINECO/FEDER, EU) and P12-FQM-1492 (Junta de Andalucía).*

## References

- [1] J. M. Arrieta, A. Rodríguez-Bernal and J. Valero, Dynamics of a reaction-diffusion equation with a discontinuous nonlinearity, *International Journal of Bifurcation and Chaos* (10), **16** (2006), 2965-2984.

# Global connections near a triple-zero singularity

A. ALGABA, M.C. DOMÍNGUEZ-MORENO, M. MERINO

*Departamento de Ciencias Integradas, Centro de Investigación de Física Teórica y Matemática FIMAT, Universidad de Huelva, Avenida de las Fuerzas Armadas s/n, 21071-Huelva (Spain)*  
algaba@dmат.uhu.es, merino@dmат.uhu.es, mcinta.dominguez@dmат.uhu.es

A.J. RODRÍGUEZ-LUIS

*Departamento de Matemática Aplicada II, E.S. Ingenieros, Universidad de Sevilla, Camino de los Descubrimientos s/n, 41092-Sevilla (Spain).*  
ajrluis@us.es

## ABSTRACT

In this work, we consider an unfolding of a normal form of the Lorenz system [1] near a triple-zero singularity

$$\begin{cases} \dot{x} &= y, \\ \dot{y} &= \epsilon_1 x + \epsilon_2 y + xz + Byz, \\ \dot{z} &= \epsilon_3 z + x^2 + Dz^2, \end{cases} \quad (1)$$

that appears when  $\epsilon_1 = \epsilon_2 = \epsilon_3 = 0$ . From this codimension-three point a double-zero bifurcation emerges when  $\epsilon_1 = \epsilon_3 = 0$  and  $\epsilon_2 \neq 0$ .

The analysis of this unfolding provides partial results that can be extended by means of the adequate numerical continuation methods [2]. Specifically, it is demonstrated that a curve of heteroclinic connections (joining the equilibria located on the  $z$ -axis) emerges from this double-zero bifurcation. Moreover, this curve has a degenerate point from which infinitely many homoclinic connections (related to one of these equilibria) emerge.

In this way, we can partially understand the dynamics in the vicinity of the triple-zero singularity exhibited by system (1).

## References

- [1] Lorenz, E.N. Deterministic non-periodic flows. *J. Atmospheric. Sci.* 20 (1963) 130-141.
- [2] Doedel, E.J., Champneys, A.R., Dercole, F., Fairgrieve, T., Kuznetsov, Y., Oldeman, B., Paffenroth, R., Sandstede, B., Wang, X., Zhang, C. AUTO-07P: Continuation and bifurcation software for ordinary differential equations (with HomCont). Technical report, Concordia University, 2012.

# Computational methods for compactons in the Rosenau–Hyman equation

FRANCISCO RUS AND FRANCISCO R. VILLATORO

*Escuela de Ingenierías Industriales, Dept. de Lenguajes y Ciencias de la Computación, Universidad de Málaga, 29071 Málaga, Spain*  
fdrus@uma.es, frvillatoro@uma.es

## ABSTRACT

Compactons are solitary waves of compact support of evolution equations with nonlinear dispersion characterized by robustness under mutual interactions [1]. The lack of analytical techniques to study the interaction of these solitary waves requires the use of numerical methods. Padé approximations with artificial viscosity have been used with success for the Rosenau–Hyman equation [2], but they introduce artificial trailing tails into the solution; they can be removed by properly designing the artificial viscosity [3]. This computational techniques can be applied to other evolution equations with compacton solutions [4]. In this oral communication their current state of the art will be reviewed.

## References

- [1] Philip Rosenau, Alon Zilburg, "Compactons," *Journal of Physics A: Mathematical and Theoretical* (AOP 25 Apr 2018), doi: 10.1088/1751-8121/aabff5.
- [2] Francisco Rus, Francisco R. Villatoro, "Padé numerical method for the Rosenau–Hyman compacton equation," *Mathematics and Computers in Simulation* 76: 188–192 (2007).
- [3] Julio Garralón, Francisco Rus, Francisco R. Villatoro, "Removing trailing tails and delays induced by artificial dissipation in Padé numerical schemes for stable compacton collisions," *Applied Mathematics and Computation* 220: 185–192 (2013).
- [4] Alejandro Apolinar-Fernández, Juan I. Ramos, "Numerical solution of the generalized, dissipative KdV-RLW-Rosenau equation with a compact method," *Communications in Nonlinear Science and Numerical Simulation* 60: 165–183 (2018).

# On the stability of positive periodic solutions in MEMS via lower and upper solutions

DANIEL NÚÑEZ

Pontificia Universidad Javeriana-Cali Colombia  
denunez@javerianacali.edu.co

## ABSTRACT

*This talk considers the existence and linear stability of positive periodic solutions for the Nathanson's model and the Comb-drive model of Micro-Electro-Mechanical-Systems (MEMS). Our results contribute to a better understanding of these models and therefore, they add up to the set of tools to decide the range of usefulness in sensing and acting of these two models. Both models are considered without damping and having an AC-DC input voltage  $\mathcal{V}(t) > 0$  of period  $T^* > 0$ . For the Nathanson's model, we prove the existence of two  $T^*$ -periodic solutions  $0 < \varphi(t) < \psi(t)$ , under the assumption that the voltage is less than the pull-in voltage and greater than a certain positive constant depending on the temporal frequency of  $\mathcal{V}$  and physical parameters. The solution  $\varphi$  is elliptic and, if the parameter  $\lambda := \frac{\min \mathcal{V}}{\max \mathcal{V}}$  is greater than a certain critical value, the solution  $\psi$  is hyperbolic. We point out that if  $\mathcal{V}$  is greater than the pull-in voltage, there are not any  $T^*$ -periodic solutions. Under the same conditions that guarantee the existence, we prove the uniqueness of these two solutions and we classify their linear stability. We can view this existence/uniqueness/linear stability classification theorem as an extension of the known results for the autonomous case when the voltage  $\mathcal{V}$  is constant and as an extension of results in [1] for the non-damping case. The Comb-drive model is studied under a cubic stiffness with coefficient  $\alpha > 0$ , and its dynamic depends on whether  $\alpha \leq 2$  or  $\alpha > 2$ . In any case, since the trivial equilibrium (the zero solution) is elliptic for an adequate range of  $\mathcal{V}$ , then it isolates this solution suggesting the existence of positive periodic solutions. We will prove the existence of periodic positive solution. When  $0 < \alpha \leq 2$ , this periodic solution is unstable for  $\lambda$  greater than a certain critical value and  $\mathcal{V}$  below the pull in voltage. For the case  $\alpha > 2$ , similar to the Nathanson case, we prove the existence of two positive periodic solutions one of which is linearized stable and the other is unstable. The methodology uses the Lower and Upper Solution Method. Some numerical examples are provided to illustrate the results.*

## References

- [1] A. Gutiérrez and P. J. Torres, *Non-autonomous saddle-node bifurcation in a canonical electrostatic MEMS*, Int. J. Bifurc. Chaos Appl. Sci. Eng. 23(5), 1350088 (9p.) (2013)



# Finite difference methods based on time-linearization for the GRLW equation

C. M. GARCÍA-LÓPEZ AND J. I. RAMOS

*Escuela de Ingenierías Industriales  
Universidad de Málaga, 29071, Málaga, Spain  
cmgl@lcc.uma.es, jirs@lcc.uma.es*

## ABSTRACT

The generalized regularized long wave (GRLW) equation is used to model unidirectional wave propagation in fluid dynamics, turbulence, acoustics, plasma dynamics, etc., and has been studied by different authors with techniques that include Petrov-Galerkin methods [1], finite difference [2, 3, 4] and finite element methods [5], and can be written as

$$u_t + \alpha u_x + \epsilon (u^p)_x = \mu u_{xx} + \delta u_{txx}, \quad -\infty < x < \infty, \quad t > 0, \quad (1)$$

where  $u$ ,  $t$  and  $x$  denote the (nondimensional) amplitude of the wave, time and spatial coordinate, respectively, and  $\alpha \geq 0$ ,  $\epsilon \geq 0$ ,  $\mu \geq 0$ ,  $\delta \geq 0$ , and  $p \geq 2$  are nondimensional parameters that are associated with linear and nonlinear advection, viscosity, dispersion, and nonlinearity, respectively. For  $\mu=0$ , the above equation has a solitary wave solution and three conserved quantities (mass, momentum and energy) [6].

In this paper, the effects of both the initial conditions and the values of the parameters on the propagation of solitary waves are assessed by considering initial conditions corresponding to the analytical solution of Eq. (1) with  $\mu=0$ , and Gaussian and triangular functions. The equation is solved numerically by means of several finite difference methods based on time-linearization techniques.

For the inviscid and viscous GRLW equations with bell-shaped initial conditions, the results show an initial adjustment of the solution whereby the leading part undergoes steepening, increases its amplitude and may break up from the trailing part. When a balance between non-linearity and dispersion is reached, a solitary wave emerges from the break-up and propagates along a straight line towards the downstream boundary in the absence of viscosity. In the presence of viscosity, the wave trajectory is curved and the wave amplitude decreases with time. For larger viscosities, solitary waves may not form. Instead, a damped wave with a steep front whose slope decreases with time is observed. The number and speed of the solitary waves that are formed after breakup increase as the dispersion parameter decreases. The effect of the linear advective term is to push the wave toward the downstream boundary. The number of solitary waves increases and the distance between solitary waves decreases as the width of the initial conditions increases regardless of their shape.

*Keywords:* GRLW equations, finite difference methods, linearization, solitary waves

## References

- [1] T. Roshan, A Petrov-Galerkin method for solving the generalized regularized long-wave (GRLW) equation, *Comput. Math. Applicat.* **63** (2012) 943-956.
- [2] C. M. García-López and J. I. Ramos, Effects of convection on a modified GRLW equation, *Appl. Math. Comput.* **219** (2012) 4118-4132.
- [3] C. M. García-López and J. I. Ramos, Solitary waves generated by bell-shaped initial conditions in the inviscid and viscous GRLW equations, *Appl. Math. Modelling*, **39** (2015) 6645-6668.
- [4] L. Zhang, A finite difference scheme for generalized regularized long-wave equation, *Appl. Math. Comput.* **168** (2005) 962-972.
- [5] H. Zeybek and S. B. G. Karakoç, A numerical investigation of the GRLW equation using lumped Galerkin approach with cubic B-spline, *SpringerPlus*, **5:199** (2016) 17 pp.
- [6] J. I. Ramos and C.M. García-López, Solitary wave formation from the Generalized Rosenau Equation. *Math. Problems Eng.*, (2016) Article ID 4618364, 17 pp.

# Time-diffracting beams in self-focusing Kerr media: self-trapped propagation below the critical power

MIGUEL A. PORRAS

*Grupo de Sistemas Complejos, Universidad Politécnica de Madrid, Spain*  
miguelangel.porras@upm.es

## ABSTRACT

*Time-diffracting beams are a new class of spatiotemporal wave packets whose diffraction-free behavior is based on spatiotemporal coupling effects, as recently described and demonstrated experimentally [1, 2, 3, 4]. Their temporal-transversal structure reproduces the axial-transversal structure of monochromatic light beams, i. e., diffraction takes place in time and not along the longitudinal direction [2, 3].*

*We have studied the nonlinear propagation of time-diffracting beams in self-focusing (cubic) Kerr media. Although stable, two-dimensional localized structures (spatial solitons) are not supported by cubic media, time-diffracting beams are demonstrated to reshape spontaneously into a propagation-invariant, seemingly robust, nonlinear wave packet of finite power. Below the critical power for collapse, the spatiotemporal structure of the attracting wave packet is that of the self-focusing of the standard monochromatic beam of the same shape and power, i. e., self-focusing is swapped from the axial direction to time. Above the critical power (and with a suitable mechanism halting collapse), the spatiotemporal structure is also that of the halted collapse of the standard monochromatic beam.*

*The well-known, unstable Townes beam, and other spatial “solitons”, represent the limit of these wave packets when localization in time is lost. These findings may open new perspectives in the longstanding problem of synthesizing truly robust, localized waves in media with the ubiquitous Kerr nonlinearity.*

## References

- [1] H. E. Kondakci and A. F. Abouraddy, “Diffraction-free space-time light sheets,” *Nature Photonics* **11**, 733 (2017).
- [2] M. A. Porras, “Gaussian beams diffracting in time,” *Opt. Lett.* **42**, 4679 (2017).
- [3] M. A. Porras, “Time-diffracting beams: on their nature, diffraction-free propagation as needles of light, and nonlinear generation,” arXiv e-prints (2017), arXiv:1802.10475 [physics.optics]
- [4] H. E. Kondakci and A. F. Abouraddy, “Airy wave packets accelerating in space-time,” arXiv e-prints (2017), arXiv: 1711.00387v1 [physics.optics].

# A necessary condition to have Weierstrass integrability

JAUME GINÉ AND JAUME LLIBRE

*Departament de Matemàtica, Inspires Research Centre, Universitat de Lleida,  
Avda. Jaume II, 69; 25001 Lleida, Catalonia, Spain*

*gine@matematica.udl.cat*

*Departament de Matemàtiques, Universitat Autònoma de Barcelona, 08193  
Bellaterra, Barcelona, Catalonia, Spain*

*jllibre@mat.uab.cat*

## ABSTRACT

*In this talk we give a necessary condition to have Weierstrass integrability. We use the solutions  $y = f(x)$  of the differential system in the plane and its associated Weierstrass cofactors. The necessary condition can be used to detect polynomial differential systems which are not Liouville integrable.*

## References

- [1] I.A. García, J. Giné, Generalized cofactors and nonlinear superposition principles, *Appl. Math. Lett.* 16 (2003), no. 7, 1137–1141.
- [3] J. Giné, M. Grau, Weierstrass integrability of differential equations, *Appl. Math. Lett.* 23 (2010), no. 5, 523–526.
- [3] J. Giné, M. Grau, J. Llibre, On the extensions of the Darboux theory of integrability, *Nonlinearity* 26 (2013), no. 8, 2221–2229.
- [4] M.F. Singer, Liouvillian first integrals of differential equations, *Trans. Amer. Math. Soc.* **333** (1992), 673–688.

# Analytic integrability of planar systems with non null linear part

Antonio Algaba, María Díaz, Cristóbal García  
Dept. Matemáticas, Facultad de Ciencias, Univ. of Huelva, Spain.  
e-mails: `algaba@uhu.es`, `crisoba@uhu.es`, `maria.diaz@dci.uhu.es`

Jaume Giné  
Departament de Matemàtica, Inspires Research Centre,  
Universitat de Lleida, Av. Jaume II, 69, 25001, Lleida, Catalonia, Spain.  
e-mail: `gine@matematica.udl.cat`

## Abstract

Recently, in [2] solved the analytic integrability problem around a nilpotent singularity for differential systems in the plane under generic conditions. In this work we solve the remaining case completing the analytic integrability problem for such singularity.

## References

- [1] A. ALGABA, C. GARCÍA, J. GINÉ, *Analytic integrability for some degenerate planar vector fields*, J. Differential Equations **257**, no. 2, (2014), 549–565.
- [2] A. ALGABA, C. GARCÍA, J. GINÉ, *The analytic integrability problem for perturbation of non-hamiltonian quasi-homogeneous nilpotent systems*, arXiv:1805.01726 [math.DS] (04 May 2018).
- [3] F. DUMORTIER, J. LLIBRE, J. C. ARTS, *Qualitative Theory of Planar Differential Systems*, Springer (2006).
- [4] A. M. LYAPUNOV, *The General Problem of the Stability of Motion*, Taylor Francis (1992).

# Suppressing chaos in damped driven systems by incommensurate excitations

R. CHACÓN, J.J.<sup>1</sup>, J.J. MIRALLES<sup>2</sup>, J.A. MARTÍNEZ<sup>3</sup>, F. BALIBREA<sup>4</sup>

<sup>1</sup>*Departamento de Física Aplicada, Escuela de Ingenierías Industriales, Universidad de Extremadura, Apartado Postal 382, E-06006 Badajoz, Spain. and Instituto de Computación Científica Avanzada (ICCAEx), Universidad de Extremadura, E-06006 Badajoz, Spain.*

<sup>2</sup>*Departamento de Física Aplicada, Escuela de Ingenieros Industriales, Universidad de Castilla-La Mancha, E-02071 Albacete, Spain.*

<sup>3</sup>*Departamento de Ingeniería Eléctrica, Electrónica y Automática, Escuela de Ingenieros Industriales, Universidad de Castilla-La Mancha, E-02071 Albacete, Spain.*

<sup>4</sup>*Departamento de Matemáticas, Facultad de Matemáticas, Universidad de Murcia, Campus de Espinardo, E-30100 Murcia, Spain.*  
<sup>1</sup>rchacon@unex.es, <sup>2</sup>juan.miralles@albacete.uned.es, <sup>3</sup>juan.mmartinez@uclm.es,  
<sup>4</sup>balibrea@um.es

## ABSTRACT

*The possibility of suppressing chaos in dissipative nonautonomous systems by additional incommensurate chaos-suppressing excitations is theoretically demonstrated through the universal example of a perturbed Duffing oscillator by considering rational approximations to the irrational ratio  $\frac{\Omega}{\omega}$  between the chaos-suppressing and chaos-inducing frequencies. For each chosen rational approximation, analytical predictions for the suitable amplitudes and initial phases of the chaos-suppressing excitation are numerically confirmed for small amplitudes and at the predicted suitable initial phases. For the significant case  $\frac{\Omega}{\omega} = \Phi$  (golden section), we study the structural stability of the suppression scenario as the respective convergents approximate the corresponding irrational ratio. Our theory predicts and numerical simulations confirm that the values of the suitable amplitudes are rather insensitive to high order convergents, while the number and values of the suitable initial phases critically depend on each particular convergent according to the maximum survival of a relevant spatio-temporal symmetry.*

## References

- [1] R. Chacón, Control of Homoclinic Chaos by Weak Periodic Perturbations (World Scientific, London, 2005).
- [2] R. Chacón, Phil. Trans. R. Soc. A 364, 2335 (2006).
- [3] R. Chacón, Phys. Rev. E 51, 761 (1995).

- [4] R. Chacón, *Europhys. Lett.* 54, 148 (2001).
- [5] W. L. Ditto, S. N. Rauseo, and M. L. Spano, *Phys. Rev. Lett.* 65, 3211 (1990).
- [6] A. Azevedo and S. M. Rezende, *Phys. Rev. Lett.* 66, 1342 (1991).
- [7] E. R. Hunt, *Phys. Rev. Lett.* 67, 1953 (1991).
- [8] R. Roy et al., *Phys. Rev. Lett.* 68, 1259 (1992).
- [9] V. Petrov, V. Gáspár, J. Masere, and K. Showalter, *Nature (London)* 361, 240 (1993).
- [9] R. Meucci, W. Gadomski, W. Ciofini, and F. T. Arecchi, *Phys. Rev. E* 49, R2528 (1994).

# The role of discrete breathers in establishing the Fourier heat transport in $\phi^4$ chain

SERGEY V. DMITRIEV

*Institute for Metals Superplasticity Problems of RAS, Khalturin St. 39, 450001  
Ufa, Russia  
dmitriev.sergey.v@gmail.com*

DAXING XIONG

*Department of Physics, Fuzhou University, Fuzhou 350108, Fujian, China  
phyxiongdx@fzu.edu.cn*

## ABSTRACT

*Anomalous (non-Fourier) heat transport is no longer just a theoretical issue since it has been observed experimentally in a number of low-dimensional nanomaterials, such as SiGe nanowires, carbon nanotubes, and others. To understand these anomalous behaviors, exploring the microscopic origin of normal (Fourier) heat transport is a fascinating theoretical topic. However, this issue has not yet been fully understood even for one-dimensional (1D) model chains, in spite of a great amount of thorough studies done to date. From those studies, it has been widely accepted that the conservation of momentum is a key ingredient to induce anomalous heat transport, while momentum-nonconserving systems usually support normal heat transport where Fourier's law is valid. But if the nonconservation of momentum is the reason, what is the underlying microscopic mechanism for the observed normal heat transport? Here we carefully revisit a typical 1D momentum-nonconserving  $\phi^4$  model, and we present evidence that the mobile discrete breathers, or, in other words, the moving intrinsic localized modes with frequency components above the linear phonon band, can be responsible for that. Main results have been published in [1].*

## References

- [1] D. Xiong, D. Saadatmand, S.V. Dmitriev. Phys. Rev. E **96**, 042109 (2017).



# Orbital–Reversibility for vector fields. Application to center problem.

A. ALGABA, I. CHECA, C. GARCÍA

*Departamento de Ciencias Integradas, Centro de investigación de física teórica y matemática FIMAT, Universidad de Huelva, Avda. Fuerzas Armadas s/n 21071 Huelva (Spain).*

isabel.checa@dmat.uhu.es

## ABSTRACT

*We consider an analytic  $n$ -dimensional autonomous system of differential equations given by:*

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}), \quad \mathbf{x} = (x_1, \dots, x_n)^T \in \mathbb{R}^n, \quad (1)$$

*having an isolated equilibrium point located at the origin. We study the orbital-reversibility problem, which consists in determining if it admits some reversibility (possibly nonlinear) modulo  $C^\infty$ -equivalence (see [3], [5], [6]). The problem of determining if system (1) has some reversibility is considered in [1], [2], [3] and [5]. In this work, we determine if there exists a time-reparametrization such that the transformed system admits some reversibility. We determine the existence of a normal form modulo reversible terms, that is, we simplify as much as possible the non-reversible terms. This allows us to give necessary conditions for the orbital-reversibility of system (1). Taking into account these conditions, we formulate a suitable algorithm to detect orbital-reversibility. We apply the results obtained to the center problem for planar vector fields. More concretely, to planar nilpotent vector fields, see [4].*

## References

- [1] A. Algaba, E. Gamero, C. García. *The reversibility problem for quasi-homogeneous dynamical systems*. Discrete and Continuous Dynamical Systems, **33** (2013) 3225–3236.
- [2] A. Algaba, C. García, M.A. Teixeira. *Reversibility and quasi-homogeneous normal forms of vector fields*. Nonlinear Analysis: Theory, Methods and Applications, **73** (2010) 510–525.
- [3] R.L. Devaney. *Reversibility, homoclinic points, and the Hénon map*. In Dynamical systems approaches to nonlinear problems in systems and circuits (N.H. Henniker, editor), SIAM, Philadelphia, PA (1988) 3–14.
- [4] A. Gasull, J. Torregrosa. *Center problem for several differential equations via Cherkas’ method*, Journal of Mathematical Analysis and Applications, 228 (1998), pp. 322–343.

- [5] J.S.W. Lamb, J.A.G. Roberts. *Time-reversal symmetry in dynamical systems: a survey*, Physica D. Nonlinear Phenomena, 112, 1-2, (1998), pp. 1–39.
- [6] D. Montgomery, L. Zippin. *Topological Transformation Groups*. Interscience Publ., New York, 1955.

# Generalising the BDS test to check i.i.d. in high-frequency financial time series

JULIO E. SANDUBETE & LORENZO ESCOT

*Faculty of Statistical Studies, Complutense University of Madrid, Spain 28040*  
jsandube@ucm.es, escot@ucm.es

## ABSTRACT

The statistical analysis of financial time series usually have considered time series sampled at fixed time intervals, mainly daily data. Recently there are some publications about models that can fit high-frequency data, even tick-by-tick. Particularly, we focus on high-frequency-based volatility models, HEAVY models [1]. Our purpose would be generalise the use of the well-known BDS test as a residual diagnostic tool of this kind of models. Brock *et. al.*, [2] proposed the BDS statistical test as a nonparametric method for testing serial independence and identically distributed hypothesis based on the correlation integral [3] that involves reconstructing the state space from a single time serie using Taken's theorem [4]. So far this reconstruction method was apply considering only time series sampled at fixed time intervals. The reconstruction of the attractors using high-frequency time series with a variable time-lapse between each observation need several considerations [5]. In this paper, we generalise the BDS as a nonlinear test considering tick-by-tick financial time series. Additionally, we check i.i.d. hypothesis in the residuals of HEAVY models as a diagnostic tool for this kind of nonlinearity modeling.

## References

- [1] Noureldin, D., Shephard, N., Sheppard, K., *Multivariate high-frequency-based volatility (HEAVY) models* J. Appl. Econ., 2012, 27:907-933
- [2] Broock, W.A., Scheinkman, J.A., Dechert, W.D. and LeBaron, B., *A test for independence based on the correlation dimension* Econometric reviews, 1996, 15(3):197-235
- [3] Grassberger, P., Procaccia, I., *Characterization of strange attractors* Physical Review Letters, 1983, 50(5), 346
- [4] Takens, F., *Detecting strange attractors in turbulence* Lecture Notes in Mathematics, 1981, 898, Springer-Verlag
- [5] Sandubete, J.E., Escot, L., *Chaotic signals inside some high-frequency financial data* Physica A: Statistical Mechanics and its Applications, Forthcoming

# Punchets: nonlinear transport in Hamiltonian pump-ratchet hybrids

THOMAS DITTRICH, NICOLÁS MEDINA SÁNCHEZ

*Depto. de Física, Universidad Nacional de Colombia, Bogotá D.C., Colombia*  
tdittrich@unal.edu.co

## ABSTRACT

*“Punchets” are hybrids between ratchets and pumps, intended as models of metal or semiconductor surfaces irradiated by a collimated laser beam [1]. They combine a spatially periodic static potential, typically asymmetric under space inversion, with a local driving that breaks time-reversal invariance. Their crucial feature is irregular driven scattering between asymptotic regions supporting periodic (as opposed to ballistic) motion. With all binary spatio-temporal symmetries broken, scattering in punchets generally induces directed currents. We here study the underlying nonlinear transport mechanisms, from chaotic scattering on the level of individual trajectories to the parameter dependence of currents evaluated as ensemble averages, in three types of Hamiltonian models, (i) with spatially periodic potentials where only in the region of driven scattering, spatial and temporal symmetries are broken, and (ii), spatially asymmetric (ratchet) potentials with a driving that only breaks time-reversal invariance. As a more realistic model of laser-irradiated surfaces, we consider (iii), a driving in the form of a running wave confined to a compact region by a static envelope. In this case, the induced current can even run against the direction of wave propagation, drastically evidencing of its nonlinear nature. Quantizing punchets is indicated as a viable research perspective.*

## References

- [1] Thomas Dittrich, Nicolás Medina Sánchez, *J. Phys. A: Math. Theor.* **51**, 055101 (2018).

# Padé numerical schemes for the sine-Gordon equation

F. MARTIN-VERGARA, F. RUS, AND F. R. VILLATORO

*Dpto. Lenguajes y Ciencias de la Computación  
Complejo de las Ingenierías. Ampliación Campus Teatinos  
Universidad de Málaga, 29071, Málaga, (Spain)  
emails: fmarver@uma.es, rusman@lcc.uma.es, villa@lcc.uma.es*

## ABSTRACT

*The sine-Gordon equation*

$$u_{tt} - u_{xx} + \sin u = 0,$$

*where the subscripts denote partial differentiation, arises in many branches of mathematical physics [1]. In this work, this equation is solved by means of four second-order time accurate finite difference methods that use Padé approximations in space [2], [3]. The accuracy and stability of the numerical solution is studied by means of the analysis discretization error and invariants for both kink-antikink and breather solutions of the sine-Gordon equation as functions of the time steps and spatial grid sizes. The new four Padé methods are compared among them and with the most widely used scheme for the sine-Gordon equation, the Strauss-Vázquez conservative method [4]. The good accuracy and stability of the Padé methods are shown after an extensive set of simulations. In terms of computational cost, the most accurate and efficient Padé scheme is the eighth-order one, as expected since the solutions of the sine-Gordon equation are smooth and the same time-integration scheme is used for all the methods.*

## References

- [1] The soliton: A new concept in applied science. Scott, Alwyn C. and Chu, F.Y.F and McLaughlin, 1973.
- [2] Calculation of weights in finite difference formulas. Fornberg, B. SIAM Rev. 40:685-691, 1998.
- [3] Self-similar radiation from numerical Rosenau-Hyman compactons. Rus, F. and Villatoro, F.R. Journal of Computational Physics 277 440-454, 2007.
- [4] Numerical solution of a nonlinear Klein-Gordon equation, Strauss, W. and Vázquez, L. Journal of Computational Physics 28 271-278, 1978.

# Localized vibrational modes in metals and alloys

ELENA A. KORZNIKOVA

*Institute for Metals Superplasticity Problems of Russian Academy of Sciences*  
elena.a.korznikova@gmail.com

## ABSTRACT

*Intrinsic localized modes or discrete breathers are studied extensively in last three decades in different physical systems. One of the most important features of the mentioned phenomenon is the ability to concentrate considerable amounts of energy in relatively small volumes. This can play a great role in case of crystal lattices because most of condensed matter objects possess crystalline long range order structure with inevitable presence of defects. Each of those has own contribution to structure dynamics and properties. Evolution of defect structure of the material is connected with overcoming of corresponding potential barriers. This process in turn is related to the energy localization in the crystal lattice. Recently it was shown that DBs can exist in pure metals and ordered alloys contributing to the evolution of crystal structure and properties. Excitation of localized modes in crystals can be performed using anzats (initial atomic displacement function ) or localization of delocalized nonlinear modes. Effectiveness of different approaches to DB excitations and their possible role in structural evolution in various metals and ordered alloys are discussed. The research was supported by Russian Scientific Foundation, Grant No 16-12-10175*

# Long-lived vortex soliton arrays nested in flat-top nonlinear Bessel beams

MIGUEL A. PORRAS<sup>1</sup>, FRANCISCO RAMOS<sup>2</sup>, JOSÉ L. GARCÍA-RIQUELME<sup>1</sup>

<sup>1</sup> *Grupo de Sistemas Complejos, ETSIME, Universidad Politécnica de Madrid, Spain*

<sup>2</sup> *Nanophotonics Technology Center, Universitat Politècnica de València, Spain*  
miguelangel.porras@upm.es

## ABSTRACT

*Ideally, optical vortex solitons need an infinite plane wave background to exist, but in practice they are nested in a broad background beam [1, 2]. Diffraction and self-defocusing broaden the background, making the vortex solitons to decay quickly, and limiting severely the implementation of their promised applications in, for instance, particle trapping, laser material processing, or in vortex-induced waveguides [3]. We report here on nonlinear Bessel beams in the self-defocusing Kerr medium as solitary-like, diffraction-free beams with arbitrarily wide and flat intensity profiles. These properties make these beams the “natural” background beams where the nested vortex solitons are seen to survive for propagation distances that are one order of magnitude larger than in the common Gaussian or super-Gaussian beam backgrounds, and where their interaction dynamics mimics the simple, and hence predictable dynamics of vortex solitons in the uniform background [4]. We have also found that dissipation does not attenuate these flat nonlinear Bessel beams; instead it contributes to their stable propagation against perturbations. This property further prolongs the distance along which the vortex solitons subsist without appreciable decay, and remain therefore useful for their intended applications [5].*

## References

- [1] A. S. Desyatnikov, L. Torner, and Y. S. Kivshar, “Optical vortices and vortex solitons,” in E. Wolf (ed.), *Progress in Optics* **47**, North-Holland, Amsterdam, 291-391 (2005).
- [2] G. A. Swartzlander and C. T. Law “Optical vortex solitons observed in Kerr nonlinear media,” *Phys. Rev. Lett.* **69**, 2503-2506 (1992).
- [3] Y. S. Kivshar and B. Luther-Davies, “Dark optical solitons: physics and applications,” *Phys. Rep.* **298**, 81-197 (1998).
- [4] M. A. Porras, F. Ramos, “Quasi-ideal dynamics of vortex solitons embedded in flat-top nonlinear Bessel beams,” *Opt. Lett.* **42**, 3275–3278 (2017).
- [5] M. A. Porras, F. Ramos, and J. L. García-Riquelme, “Long-lived optical vortex solitons in self-defocusing dissipative media,” in preparation.

## Participants

A. Algaba. Universidad de Huelva, Spain  
F. Balibrea. Universidad de Murcia, Spain  
Rubén Caballero. Universidad Miguel Herandez de Elche, Spain  
Luis Carretero. Universidad Miguel Hernández de Elche, Spain  
Alexandre N. Carvalho. Universidade de São Paulo, Brazil  
R. Chacón. Universidad de Extremadura, Spain  
I. Checa. Universidad de Huelva, Spain  
Nathan Craig. Univeristy of Newcastle, UK  
Thomas Dittrich. Universidad Nacional de Colombia, Colombia.  
María Díaz. Universidad de Huelva, Spain  
Sergey V. Dmitriev. Ufa, Russia  
M.C. Domínguez-Moreno. Universidad de Huelva, Spain  
Angel Duran. University of Valladolid, Spain  
Lorenzo Escot. Complutense University of Madrid, Spain  
Ivan R. Garashchuk. National Research Nuclear University, Moscow  
Cristobal García. Universidad de Huelva, Spain  
Isaac A. García. Universitat de Lleida, Spain  
C. M. García-López. Universidad de Málaga, Spain  
José L. Carcía-Riquelme. Universidad Politécnica de Madrid, Spain  
Jaume Giné. Universitat de Lleida, Spain  
Harriet Grigg. Univeristy of Newcastle, UK  
Peter Jan Van Leeuwen. University of Reading, UK  
Elena A. Korznikova, Russian Academy of Sciences  
Nikolay A. Kudryashov. National Research Nuclear University, Moscow  
Jaume Llibre. Universitat Autònoma de Barcelona, Spain  
Magdalena Lucivi. University of Reading, UK  
Pedro Marín-Rubio. Universidad de Sevilla, Spain  
F. Martín-Vergara. Universidad de Málaga, Spain  
J. A. Martínez. Universidad de Castilla-La Mancha, Spain  
Susanna Maza. Universitat de Lleida, Spain  
Nicolás Medina Sánchez. Universidad Nacional de Colombia, Colombia.  
M. Merino. Universidad de Huelva, Spain  
J.J. Miralles. Universidad de Castilla-La Mancha, Spain  
Daniel Núñez. Pontificia Universidad Javeriana-Cali. Colombia  
Miguel A. Porras. Universidad Politécnica de Madrid, Spain.  
Manuel Pulido. University of Reading, UK  
Francisco Ramos. Universidad Politécnica de Valencia, Spain  
J. I. Ramos. Universidad de Málaga, Spain.  
A.J. Rodriguez-Ruiz. Universidad de Sevilla, Spain  
Santiago Rosa. Universidad Nacional de Cordoba, Argentina  
Francisco Rus. Universidad de Málaga, Spain  
Julio E. Sandubete. Complutense University of Madrid, Spain  
Cimetry I. Sinelshchikov. National Research Nuclear University, Moscow  
José Valero. Universidad Miguel Hernández de Elche, Spain  
Francisco R. Villatoro. Universidad de Málaga, Spain  
Santiago Wiedman. Universidad Nacional del Nordeste, Argentina  
Daxing Xiong. Fuzhou University, China